

Can $(dG/dt)/G$ Bound the Local Cosmological Dynamics?

Yurii V. Dumin*

Theoretical Department, IZMIRAN, Russian Academy of Sciences, Troitsk, Moscow reg., 142190 Russia

(Dated: October 9, 2006)

It is argued that constraints on time variation of the gravitational constant $(dG/dt)/G$, e.g. derived from the lunar laser ranging, cannot be immediately applied to restrict the cosmological expansion at planetary scales, as it was done by Williams, Turyshev, and Boggs [PRL, **93**, 261101 (2004)].

Paper [1] presented the latest results of the analysis of lunar laser ranging (LLR), one of which is a constraint on time variation in the gravitational constant as strong as $\dot{G}/G = (4 \pm 9) \times 10^{-13} \text{ yr}^{-1}$. So, as was concluded by these authors, “the $\dot{G}/G$ uncertainty is 83 times smaller than the inverse age of the Universe, $t_0 = 13.4 \text{ Gyr} \dots$ Any isotropic expansion of the Earth’s orbit which conserves angular momentum will mimic the effect of $\dot{G}$ on the Earth’s semimajor axis, $\dot{a}/a = -\dot{G}/G \dots$ There is no evidence for such local ($\sim 1 \text{ AU}$) scale expansion of the solar system.”

Unfortunately, the above-stated equivalence between the effect of variable G and the cosmological expansion is based solely on the Newtonian arguments. A more accurate treatment of this problem in the framework of General Relativity for a general case of the multi-component Universe is very difficult. Nevertheless, for the particular case of a point-like mass M in the Universe filled only with Λ -term (or the so-called ‘dark energy’, which is now assumed to be the main ingredient responsible for the cosmological dynamics), the corresponding solution was obtained very long time ago by Kottler [2]:

$$ds^2 = - \left(1 - \frac{2GM}{c^2 r'} - \frac{\Lambda r'^2}{3} \right) c^2 dt'^2 + \left(1 - \frac{2GM}{c^2 r'} - \frac{\Lambda r'^2}{3} \right)^{-1} dr'^2 + r'^2 (d\theta^2 + \sin^2 \theta d\varphi^2), \quad (1)$$

where G is the gravitational constant, and c is the speed of light. (For a general review, see also [3].)

After a transformation to the cosmological Robertson–Walker coordinates

$$r' = a_0 \exp\left(\frac{ct}{r_0}\right) r, \quad (2)$$

$$t' = t - \frac{1}{2} \frac{r_0}{c} \ln \left[1 - \frac{a_0^2}{r_0^2} \exp\left(\frac{2ct}{r_0}\right) r^2 \right], \quad (3)$$

metric (1) takes the form

$$ds^2 = g_{tt} c^2 dt^2 + 2 g_{tr} c dt dr + g_{rr} dr^2 + g_{\theta\theta} d\theta^2 + g_{\varphi\varphi} d\varphi^2, \quad (4)$$

where

$$g_{tt} = \frac{- \left(1 - \frac{r_g}{r'} - \frac{r'^2}{r_0^2} \right)^2 + \left(1 - \frac{r'^2}{r_0^2} \right)^2 \frac{r'^2}{r_0^2}}{\left(1 - \frac{r_g}{r'} - \frac{r'^2}{r_0^2} \right) \left(1 - \frac{r'^2}{r_0^2} \right)}, \quad (5)$$

$$g_{tr} = \frac{\left(1 - \frac{r'^2}{r_0^2} \right)^2 - \left(1 - \frac{r_g}{r'} - \frac{r'^2}{r_0^2} \right)^2 \frac{r'}{r_0}}{\left(1 - \frac{r_g}{r'} - \frac{r'^2}{r_0^2} \right) \left(1 - \frac{r'^2}{r_0^2} \right)} \frac{r'}{r}, \quad (6)$$

$$g_{rr} = \frac{\left(1 - \frac{r'^2}{r_0^2} \right)^2 - \left(1 - \frac{r_g}{r'} - \frac{r'^2}{r_0^2} \right)^2 \frac{r'^2}{r_0^2}}{\left(1 - \frac{r_g}{r'} - \frac{r'^2}{r_0^2} \right) \left(1 - \frac{r'^2}{r_0^2} \right)} \frac{r'^2}{r^2}, \quad (7)$$

$$g_{\theta\theta} = g_{\varphi\varphi} / \sin^2 \theta = r'^2. \quad (8)$$

In the above formulas, $r_g = 2GM/c^2$, $r_0 = \sqrt{3/\Lambda}$, and a_0 is the scale factor of FRW Universe.

Taking $a_0 = 1$ at $t = 0$ and keeping only the lowest-order terms of r_g and $1/r_0$, we get

$$g_{tt} \approx - \left[1 - \frac{2GM}{c^2 r} \left(1 - \frac{c\sqrt{\Lambda} t}{\sqrt{3}} \right) \right], \quad (9)$$

$$g_{tr} \approx \frac{4GM\sqrt{\Lambda}}{\sqrt{3} c^2}, \quad (10)$$

$$g_{rr} \approx \left[1 + \frac{2GM}{c^2 r} \left(1 - \frac{c\sqrt{\Lambda} t}{\sqrt{3}} \right) \right] \left(1 + \frac{2c\sqrt{\Lambda} t}{\sqrt{3}} \right), \quad (11)$$

$$g_{\theta\theta} = g_{\varphi\varphi} / \sin^2 \theta \approx r^2 \left(1 + \frac{2c\sqrt{\Lambda} t}{\sqrt{3}} \right). \quad (12)$$

As is seen from the above expressions, manifestation of the Λ -term in some components of the metric tensor really looks like the influence of variable G if we assume that $G = G_0 + \dot{G}t$, where $\dot{G} = -c\sqrt{\Lambda/3}$. Unfortunately, such interpretation is not self-consistent: the Λ -dependence of a few other components is irreducible to the variable coefficient of gravitational coupling.

Therefore, the available limits on $\dot{G}/G$, in general, cannot be reinterpreted as a constraint on the local cosmological dynamics.

* dumin@yahoo.com

- [1] J. G. Williams, S. G. Turyshev, and D. H. Boggs, Phys. Rev. Lett. **93**, 261101 (2004).
- [2] F. Kottler, Ann. Phys. (Leipzig) **56**, 401 (1918).
- [3] D. Kramer, H. Stephani, M. MacCallum, and E. Herlt, *Exact Solutions of the Einsteins Field Equations* (Deutscher Verlag der Wissenschaften, Berlin, 1980).